

**Algebra Qualifying Exam**  
**January 2007**

Do all 5 problems.

1. Let  $G$  be a finite group and let  $\text{Syl}_p(G)$  denote its set of Sylow  $p$ -subgroups.
  - a. Suppose that  $S$  and  $T$  are distinct members of  $\text{Syl}_p(G)$  chosen so that  $S \cap T$  is maximal among all such intersections. Prove that the normalizer  $\mathbf{N}_G(S \cap T)$  has more than one Sylow  $p$ -subgroup. (5 points)
  - b. Show that  $S \cap T = 1$  for all  $S, T \in \text{Syl}_p(G)$ , with  $T \neq S$ , if and only if  $\mathbf{N}_G(P)$  has exactly one Sylow  $p$ -subgroup for every nonidentity  $p$ -subgroup  $P$  of  $G$ . (5 points)
2. Let  $R$  be a commutative, Noetherian integral domain.
  - a. If  $P$  is a prime ideal of  $R$ , show that the radical of  $P^n$  is  $P$ . (2 points)
  - b. If  $R$  has a unique nonzero prime ideal  $P$ , prove that all ideals of  $R$  are primary. (3 points)
  - c. Conversely, let us now assume that all ideals of  $R$  are primary, and let  $P$  and  $Q$  be distinct prime ideals of  $R$  with  $Q \not\supseteq P$ . Since  $P^n \cap Q$  is primary, deduce first that  $P^n \supseteq Q$  and then that  $Q = 0$ . (Hint. Consider whether the intersection  $P^n \cap Q$  can be irredundant.) (5 points)
3. Let  $F$  be a field of characteristic 0 and let  $f \in F[X]$  be an irreducible polynomial of degree  $> 1$  with splitting field  $E \supseteq F$ . Define  $\Omega = \{\alpha \in E \mid f(\alpha) = 0\}$ .
  - a. Let  $\alpha \in \Omega$  and let  $m$  be a positive integer. If  $g \in F[X]$  is the minimal polynomial of  $\alpha^m$  over  $F$ , show that  $\{\beta^m \mid \beta \in \Omega\}$  is the set of roots of  $g$ . (3 points)
  - b. Now fix  $\alpha \in \Omega$  and suppose that  $\alpha r \in \Omega$  for some  $r \in F$ . Show that, for all  $\beta \in \Omega$  and integers  $i \geq 0$ , we have  $\beta r^i \in \Omega$ . Conclude that  $r$  is a root of unity. (3 points)
  - c. If  $\alpha$  and  $r$  are as in (b) and if  $m$  is the multiplicative order of the root of unity  $r$ , show that  $f(X) = g(X^m)$ , where  $g$  is the minimal polynomial of  $\alpha^m$  over  $F$ . (4 points)
4. Let  $V$  be a finite dimensional vector space over a field  $K$  and assume that  $V$  is endowed with a not necessarily symmetric bilinear form  $\langle \cdot, \cdot \rangle: V \times V \rightarrow K$ . We let  $R$  and  $L$  denote the right and left radicals of  $\langle \cdot, \cdot \rangle$  given by  $R = \{x \in V \mid \langle V, x \rangle = 0\}$  and  $L = \{x \in V \mid \langle x, V \rangle = 0\}$ , so that these are both subspaces of  $V$ .
  - a. Use the bilinear form to construct a linear transformation  $T$  from  $V$  to the dual space  $(V/R)^*$  of  $V/R$  such that  $\ker(T) = L$ . (6 points)
  - b. Show that  $\dim_K L = \dim_K R$ , and deduce that the map  $T$  is surjective. (4 points)
5. Let  $A$  be an additive abelian group and let  $B$  be a subgroup. We say that  $B$  is essential in  $A$ , and write  $B \text{ ess } A$ , if and only if  $B \cap X \neq 0$  for all nonzero subgroups  $X$  of  $A$ .
  - a. If  $B_1 \text{ ess } A_1$  and  $B_2 \text{ ess } A_2$ , prove that  $(B_1 \oplus B_2) \text{ ess } (A_1 \oplus A_2)$ . (5 points)
  - b. If  $B \text{ ess } A$ , and  $B$  has no nonzero elements of finite order, prove that  $A$  has no nonzero elements of finite order. (2 points)
  - c. Let  $Q$  denote the additive group of rational numbers and suppose that  $Q \text{ ess } A$ , for some abelian group  $A$ . Prove that  $Q = A$ . (3 points)